

VIRTUAL UNIVERSITY OF PAKISTAN

A Good Education is a Foundation For a Better Future

CORRECT ANSWER SOLVED BY HADI
EMAIL: hadirajputofficial@gmail.com
WHATSAPP& CALL: 03087122922

TEAM HADI(VU LAHORE)

MID TERM PAST PAPER

We Deal All Kind of VU Project
(PHP, .NET, ANDROID, PYTHON)

MADE AND SOLVED BY HADI

WARNING: Team HADI is not responsible for any mistake or wrong answer. All students reading and using this document may check and confirm the answers at their own.



WhatsApp: 03087122922



FACEBOOK ID: <https://www.facebook.com/hadipastpapers/>

Best of luck!



Question No : 1 of 26

Marks: 1 (Budgeted Time 1 Min)

The form of the exact solution to

$$2\frac{dy}{dx} + 3y = e^{-x}, y(0) = 5$$

is

Answer (Please select your correct option)

☐ $Ae^{-1.5x} + Bxe^{-x}$

☐ $Ae^{1.5x} + Be^{-x}$

☐ $Ae^{1.5x} + Bxe^{-x}$

☐ $Ae^{-1.5x} + Be^{-x}$



correct answer solve
by hadi

MADE BY HADI

Email: hadirajputofficial@gmail.com Whatsapp: 0308 7122 922 (Team Hadi L


Question No : 2 of 26

Marks: 1 (Budgeted Time 1 Min)

The differential equation $2\frac{dy}{dx} + x^2y = 2x + 3, y(0) = 5$ is

Answer (Please select your correct option)

☐ Linear



correct answer solve
by hadi

☐ Nonlinear

☐ Linear with fixed constants

☐ Undeterminable to be linear or nonlinear

MADE BY HADI

Email: hadirajputofficial@gmail.com Whatsapp: 0308 7122 922 (Team Hadi Lahore)

Question No : 3 of 26

Marks: 1 (Budgeted Time 1 Min)

The value of amplitude in the solution $X=40\sin(7t+4)$ is


Answer (Please select your correct option)

☐ 7

☐ 4

☒ 40

☐ 20



correct answer solve
by hadi

MADE BY HADI

Question No : 4 of 26

Marks: 1 (Budgeted Time 1 Min)

The order of the differential equation $(\frac{\partial^2 u}{\partial t^2})^3 + \frac{\partial u}{\partial t} = 0$ is


Answer (Please select your correct option)

☐ 0

☐ 1

☒ 2

☐ 3



correct answer solve
by hadi

MADE BY HADI


Question No : 5 of 26

Marks: 1 (Budgeted Time 1 Min)

Which of the following is the linear differential equation?

Answer (Please select your correct option)

☐ $x \frac{dy}{dx} + y = 0$



correct answer solve
by hadi

☐ $y \frac{dy}{dx} + x = 0$

☐ $x \frac{d^2y}{dx^2} + \frac{dy}{dx} + y^2 = 0$

☐ $\frac{dy}{dx} = e^y$

MADE BY HADI

Email: hadirajputofficial@gmail.com Whatsapp: 03087122922 (Team Hadi Lahore)

Question No : 6 of 26

Marks: 1 (Budgeted Time 1 Min)

A solution obtained by giving particular values to the arbitrary constants in the General Solution of a differential equation is called a -----


Answer (Please select your correct option)

☐ Singular solution

☒ Particular solution

☐ Explicit Solution

☐ None of these



correct answer solve
by hadi

MADE BY HADI

Email: hadirajputofficial@gmail.com Whatsapp: 03087122922 (Team Hadi Lahore)

Question No : 7 of 26


Marks: 1 (Budgeted Time 1 Min)

Which of the following equations satisfy the differential equation $\frac{dy}{dx} = x$

Answer (Please select your correct option)

☐

$$2y = x^2 + c$$



correct answer solve
by hadi

☐

$$y = x^2 + c$$

☐

$$y = x^2$$

☐

$$y = x + c$$

MADE BY HADI

Email: hadirajputofficial@gmail.com Whatsapp: 0308 7122 922 (Team Hadi Lahore)

Question No : 8 of 26

Marks: 1 (Budgeted Time 1 Min)

A differential equation $\frac{dy}{dx} + p(x)y = q(x)y^n$ for $n \neq 0, 1$ is called a/an ----.

Answer (Please select your correct option)

☐ Non-Exact differential equation

☐ Linear differential equation

☒ Bernoulli equation

☐ Exact differential equation



correct answer solve
by hadi

MADE BY HADI


Question No : 9 of 26

Marks: 1 (Budgeted Time 1 Min)

In a Bernoulli equation $\frac{dy}{dx} + \frac{1}{x}y = xy^2$, identify $p(x)$, $q(x)$ & n respectively.

Answer (Please select your correct option)

☐ $\frac{1}{x}, x \text{ \& } 2$



correct answer solve
by hadi

☐ $y, x^2y \text{ \& } 2$

☐ $y, \frac{1}{x} \text{ \& } 2$

☐ $\frac{1}{y}, y \text{ \& } 2$

MADE BY HADI

Email: hadirajputofficial@gmail.com Whatsapp: 0308 7122 922 (Team Hadi Lahore)

Question No : 10 of 26

Marks: 1 (Budgeted Time 1 Min)

In order to change the Bernoulli Equation

$$\frac{dy}{dx} + p(x)y = q(x)y^n$$

into linear differential equation, we choose -----


Answer (Please select your correct option)

☐ $v = y^{n-1}$

☒ $v = y^{1-n}$

☐ $v = y^n$

☐ $v = y'$



correct answer solve
by hadi

MADE BY HADI

Email: hadirajputofficial@gmail.com Whatsapp: 03087122922 (Team Hadi Lahore)

Question No : 11 of 26


Marks: 1 (Budgeted Time 1 Min)

The differential equation ----- is separable.

Answer (Please select your correct option)

☐ $x(x+y) \frac{dy}{dx} = 4$

☐ $\frac{dy}{dx} = \frac{x^2}{x+xy}$



correct answer solve
by hadi

☐ $\frac{dy}{dx} = \frac{y}{1+xy^3}$

☐ $\frac{dy}{dx} = \frac{xy+3}{1+2xy}$

MADE BY HADI

Email: hadirajputofficial@gmail.com Whatsapp: 03087122922 (Team Hadi Lahore)

Question No : 12 of 26

Marks: 1 (Budgeted Time 1 Min)

If $y = x$, then $\frac{dy}{dx} = \dots\dots\dots$

Answer (Please select your correct option)

☐ 0

☒ 1

☐ x

☐ $x+1$



correct answer solve
by hadi

MADE BY HADI

Email: hadirajputofficial@gmail.com Whatsapp: 0308 7122 922 (Team Hadi Lahore)

Question No : 13 of 26

Marks: 1 (Budgeted Time 1 Min)

If $f(x)$ and $g(x)$ are linearly dependent on I , then-----

Answer (Please select your correct option)

- ☐ $W(f,g)(x) \neq 0$ (Wronskian) for all x in the interval I .
- ☐ $W(f,g)(x) = 0$ (Wronskian) for all x in the interval I .
- ☐ $W(f,g)(x)$ may or may not be zero for all x in the interval I .
- ☐ $W(f,g)(x)$ is not defined for all x in the interval I .



correct answer solve
by hadi

MADE BY HADI

Email: hadirajputofficial@gmail.com Whatsapp: 0308 7122 922 (Team Hadi Lahore)

Question No : 14 of 26

Marks: 1 (Budgeted Time 1 Min)

Wronskian of $\sin x, \cos x$ is----


Answer (Please select your correct option)

☐ $\sin x$

☐ 1

☒ -1

☐ $\cos x$



MADE BY HADI

Email: hadirajputofficial@gmail.com Whatsapp: 0308 7122 922 (Team Hadi Lahore)

Question No : 15 of 26

Marks: 1 (Budgeted Time 1 Min)

The orthogonal trajectory to the family of curves $3x + 4y = c$ is-----.


Answer (Please select your correct option)

☐ $y = 2x$

☐ $y - x = c$

☒ $4x - 3y = c$

☐ $4x + 3y = c$




MADE BY HADI

Email: hadirajputofficial@gmail.com Whatsapp: 0308 7122 922 (Team Hadi Lahore)

If T denotes the half life and the initial amount of the substance be A_0 , then the amount of radioactive substance at time T is given by-----.

Answer (Please select your correct option)

☐ $A(T) = \frac{A_0}{2}$



correct answer solve
by hadi

☐ $A(T) = \frac{T}{2}$

☐ $A(T) = A_0$

☐ $A(T) = \frac{kT}{2}$

MADE BY HADI

Email: hadirajputofficial@gmail.com Whatsapp: 0308 7122 922 (Team Hadi Lahore)

Question No : 17 of 26

Marks: 1 (Budgeted Time 1 Min)

If the auxiliary equation has roots $-2, -2$, then the general solution of the differential equation is _____


Answer (Please select your correct option)

☐ $y = c_1 e^{-2x} + c_2 e^{-2x}$

☐ $y = c_1 e^{-2+2x}$

☐ $y = c_1 e^{2x} + c_2 e^{-2x}$

☐ $y = (c_1 + c_2 x) e^{-2x}$



correct answer solve
by hadi


MADE BY HADI

If the auxiliary equation has complex roots $-2 \pm 3i$, then the general solution of the differential equation is _____

Answer (Please select your correct option)

☐ $y = ce^{2x}(\cos 10x + \sin 10x)$

☐ $y = e^{-2x}(c_1 \cos 3x + c_2 \sin 3x)$



correct answer solve
by hadi

☐ $y = e^{ix}(c_1 \cos 2x + c_2 \sin 2x)$

☐ $y = e^{2x}(c_1 \cos x - c_2 \sin x)$

MADE BY HADI

Email: hadirajputofficial@gmail.com Whatsapp: 03087122922 (Team Hadi Lahore)

If $y = e^{-6x}$, then which of the following is the most accurate option?

Answer (Please select your correct option)

☐ The annihilator operator of $y = e^{-6x}$ is $(D - 6)^2$.

☐ The annihilator operator of $y = e^{-6x}$ is $(D - 6)^3$.

☐ The annihilator operator of $y = e^{-6x}$ is $(D - 6)$.

☐ The annihilator operator of $y = e^{-6x}$ is $(D + 6)$.




correct answer solve
by hadi

MADE BY HADI

If the complementary solution of $\frac{d^2y}{dx^2} - \frac{dy}{dx} - 2y = 3\sin x$ is $c_1 e^{-x} + c_2 e^{2x}$, then which of the following is the most accurate option for it?

Answer (Please select your correct option)

- ☐ The general form of the particular solution will be $(A \sin x + B \cos x) e^x$.
- ☐ The general form of the particular solution will be $A x \sin x + B x \cos x$.
- ☐ The general form of the particular solution will be $(A x \sin x + B x \cos x) e^{-x}$.
- ☐ The general form of the particular solution will be $A \sin x + B \cos x$.



correct answer solve
by hadi

MADE BY HADI

Email: hadirajputofficial@gmail.com Whatsapp: 0308 7122922 (Team Hadi Lahore)

Evaluate $e^{-\int P(x) dx}$ using the differential equation $x^2 \frac{d^2 y}{dx^2} + x \frac{dy}{dx} + \left(x^2 - \frac{1}{4}\right)y = 0$ by writing it in the standard form $\frac{d^2 y}{dx^2} + P(x) \frac{dy}{dx} + Q(x)y = 0$.

Answer ([Please click here to Add Answer](#))



MADE BY HADI

Email: hadirajputofficial@gmail.com Whatsapp: 03087122922 (Team Hadi Lahore)

If the complementary solution of the following differential equation is $c_1 \sin 2x + c_2 \cos 2x$, then what will be the general form of its particular solution:

$$\frac{d^2 y}{dx^2} + 4y = \cos 2x$$

Answer ([Please click here to Add Answer](#))



MADE BY HADI

Email: hadirajputofficial@gmail.com Whatsapp: 03087122922 (Team Hadi Lahore)

Question No : 23 of 26

Marks: 3 (Budgeted Time 6 Min)

If a mass weighing $8lb$ and $k = 40 lb/ft$ then find the amount of elongation's.

Answer ([Please click here to Add Answer](#))



MADE BY HADI

Email: hadirajputofficial@gmail.com Whatsapp: 0308 7122 922 (Team Hadi Lahore)

If $y_1(x) = e^{-4x}$ is the first solution of $\frac{d^2 y}{dx^2} + 6 \frac{dy}{dx} + 8y = 0$, then construct its second solution using $y_2(x) = y_1(x) \int \frac{e^{-\int P(x) dx}}{y_1^2(x)} dx$.

Note: where $P(x)$ is a coefficient of the first derivative involving in the above given differential equation.

Answer ([Please click here to Add Answer](#))



MADE BY HADI

Email: hadirajputofficial@gmail.com Whatsapp: 0308 7122 922 (Team Hadi Lahore)

Question No : 25 of 26

Marks: 5 (Budgeted Time 10 Min)

A mass weighing $2lb$ stretches a spring 6 inches. Determine the equation of simple harmonic motion.

Answer ([Please click here to Add Answer](#))



MADE BY HADI

Email: hadirajputofficial@gmail.com Whatsapp: 03087122922 (Team Hadi Lahore)

Question No : 26 of 26

Marks: 5 (Budgeted Time 10 Min)

A radioactive isotope has a half-life of 16 days. We have 45 g at the end of 32 days. How much radioisotope was initially present?

Answer ([Please click here to Add Answer](#))



MADE BY HADI

Email: hadirajputofficial@gmail.com Whatsapp: 0308 7122 922 (Team Hadi Lahore)

HADI VU PAST PAPER

WE DEAL ALL KIND OF VU PROJECT
(PHP, ANDROID, PYTHON)

DEAR STUDENTS

ARE YOU LOOKING
FOR ASSISTANCE

- 1) ASSIGNMENTS
- 2) QUIZZES
- 3) GDB
- 4) GRAND QUIZZES

We Provide, You Solution
of VU Academic Activities
include Assignments,
Quizzes, GDBs, Grand Quizzes
If You Need Help in Services
Kindly inform us.

TEAM HADI
(VU LAHORE)



TEAM HADI
(VU LAHORE)



HADI TECH

#PAID_TASK

CONTACT US

CALL & WHATSAPP

0308-7122922